

Additional Exercises 2.3
Form I
Solving Linear Equations

Solve the equation.

1. $7x - (3x - 1) = 2$ 1. _____

2. $6(4x - 1) = 24$ 2. _____

3. $(y - 2) - (y + 8) = 4y$ 3. _____

4. $3(5x - 2) = 9$ 4. _____

5. $2(2 + 4x) = 52$ 5. _____

6. $-2(2x - 1) = -4$ 6. _____

7. $1.5x + 0.9 = 1.2x - 1.5$ 7. _____

8. $1.4x - 1.1 = 0.2x + 1.3$ 8. _____

Solve each equation. Begin your work by rewriting each equation without fractions.

9. $\frac{f}{5} - 4 = 1$ 9. _____

10. $\frac{2x}{5} - \frac{x}{3} = 3$ 10. _____

11. $\frac{x}{7} = 10 - \frac{x}{3}$ 11. _____

12. $\frac{x}{3} - 1 = -\frac{x}{2}$ 12. _____

13. $\frac{3x}{10} - \frac{1}{2} = \frac{x}{5}$ 13. _____

14. $\frac{x}{2} = \frac{x+5}{3}$ 14. _____

15. $\frac{2x}{3} + \frac{5}{6} = \frac{3x}{2}$ 15. _____

Name _____

Date _____

16. Forensic scientists use the length of certain bones to calculate the height of a person. When the femur (f), the bone from the knee to the hip socket is used, the following formula applies for men: $h = 69.09 + 2.24f$, where h is the height and f is the length of the femur. Find the height of a man with a femur measuring 59 centimeters. 16. _____
17. There is a formula that gives a correspondence between women's shoe sizes in the United States and those in Italy. The formula is $S = 2(x + 12)$, where S is the size in Italy and x is the size in the United States. What would be the US size for an Italian size of 32? 17. _____
18. In one state, speeding fines are determined by the formula $F = 6(x - 60) + 75$, where F is the cost, in dollars, of the fine if a person is caught driving x miles per hour. If the fine comes to \$249, how fast was the person driving? 18. _____
19. When you buy an item on which sales tax is charged, the total cost is calculated by the formula $T = P + \frac{S}{100}P$, where T is the total cost, P is the item's price, and S is the sales tax rate (as a percent). If you pay \$20.045 for an item priced at \$19, what is the tax rate? 19. _____
20. To convert a Fahrenheit temperature to Celsius, one formula to use is $F = \frac{9}{5}C + 32$ where F is the Fahrenheit temperature (in degrees) and C is the Celsius temperature. What is the Celsius temperature (to the nearest degree) when Fahrenheit temperature is 86° ? 20. _____

Additional Exercises 2.3
Form II
Solving Linear Equations

Solve the equation.

1. $9(x + 2) = 3(x - 2)$ 1. _____

2. $7(5x - 2) = 6(6x - 1)$ 2. _____

3. $2(x + 1) = 29 - x$ 3. _____

4. $13(6x - 7) = 8x - 2$ 4. _____

5. $4(x + 3) = 5(x - 7)$ 5. _____

6. $5(3 + x) - x = 4(x + 2) + 7$ 6. _____

7. $0.7x - 2.1 = 1.5x + 2.7$ 7. _____

8. $0.2x - 1.2 = 1.4x - 3.6$ 8. _____

Solve each equation. Begin your work by rewriting each equation without fractions.

9. $\frac{x}{3} + \frac{x}{4} = \frac{7}{4}$ 9. _____

10. $\frac{3x}{5} = \frac{1}{20} + \frac{x}{2}$ 10. _____

11. $\frac{x}{5} - \frac{1}{5} = -3$ 11. _____

12. $\frac{x}{7} - 7 = -3$ 12. _____

13. $\frac{2x}{5} = \frac{x}{3} + 5$ 13. _____

14. $3x - 1 = \frac{x}{5} + \frac{4x}{5}$ 14. _____

15. $\frac{x+3}{3} - \frac{x}{4} = \frac{x-2}{5}$ 15. _____

Name _____

Date _____

16. Forensic scientists use the length of certain bones to calculate the height of a person. When the femur (f), the bone from the knee to the hip socket is used, the following formula applies for men: $h = 69.09 + 2.24f$, where h is the height and f is the length of the femur. Find the height of a man with a femur measuring 57 centimeters. 16. _____
17. There is a formula that gives a correspondence between women's shoe sizes in the United States and those in Italy. The formula is $S = 2(x + 12)$, where S is the size in Italy and x is the size in the United States. What would be the US size for an Italian size of 30? 17. _____
18. In one state, speeding fines are determined by the formula $F = 6(x - 60) + 75$, where F is the cost, in dollars, of the fine if a person is caught driving x miles per hour. If the fine comes to \$225, how fast was the person driving? 18. _____
19. When you buy an item on which sales tax is charged, the total cost is calculated by the formula $T = P + \frac{S}{100}P$, where T is the total cost, P is the item's price, and S is the sales tax rate (as a percent). If you pay \$20.235 for an item priced at \$19, what is the tax rate? 19. _____
20. To convert a Fahrenheit temperature to Celsius, one formula to use is $F = \frac{9}{5}C + 32$ where F is the Fahrenheit temperature (in degrees) and C is the Celsius temperature. What is the Celsius temperature (to the nearest degree) when Fahrenheit temperature is 95° ? 20. _____

Additional Exercises 2.3
Form III
Solving Linear Equations

Solve the equation.

1. $7(x-3) - 2x = 5(x-3)$ 1. _____

2. $3(x-4) = 6(x-3)$ 2. _____

3. $3(x+2) - 4x = x+16$ 3. _____

4. $5x + 2(1-x) = 2(2x-1)$ 4. _____

5. $4(x+2) = 14 - 2(3-2x)$ 5. _____

6. $5(x+3) + 9 = 3(x-2) + 6$ 6. _____

7. $1.7x + 2.8 = 1.3x - 3.6$ 7. _____

8. $3.1x + 0.5 = 2.7x + 1.7$ 8. _____

Solve each equation. Begin your work by rewriting each equation without fractions.

9. $\frac{1}{2}x + 2 - \frac{2}{3}x + \frac{2}{3} = 3$ 9. _____

10. $\frac{1}{4}x - 1 = \frac{1}{2}x + \frac{3}{2}$ 10. _____

11. $2x - \frac{10}{3} = 6 - \frac{1}{3}x$ 11. _____

12. $\frac{1}{4}x - \frac{1}{12} = \frac{1}{6}x + \frac{1}{6}$ 12. _____

13. $\frac{5}{2} = \frac{3}{2}x + \frac{7}{4}$ 13. _____

14. $\frac{r}{5} + \frac{6}{5} = \frac{r}{7} + \frac{8}{7}$ 14. _____

15. $\frac{y}{5} - \frac{2}{5} = \frac{1}{3} - y$ 15. _____

Name _____

Date _____

16. Forensic scientists use the length of certain bones to calculate the height of a person. When the femur (f), the bone from the knee to the hip socket is used, the following formula applies for men: $h = 69.09 + 2.24f$, where h is the height and f is the length of the femur. Find the height of a man with a femur measuring 53 centimeters. 16. _____
17. There is a formula that gives a correspondence between women's shoe sizes in the United States and those in Italy. The formula is $S = 2(x + 12)$, where S is the size in Italy and x is the size in the United States. What would be the US size for an Italian size of 34? 17. _____
18. In one state, speeding fines are determined by the formula $F = 6(x - 60) + 75$, where F is the cost, in dollars, of the fine if a person is caught driving x miles per hour. If the fine comes to \$261, how fast was the person driving? 18. _____
19. When you buy an item on which sales tax is charged, the total cost is calculated by the formula $T = P + \frac{S}{100}P$, where T is the total cost, P is the item's price, and S is the sales tax rate (as a percent). If you pay \$20.425 for an item priced at \$19, what is the tax rate? 19. _____
20. To convert a Fahrenheit temperature to Celsius, one formula to use is $F = \frac{9}{5}C + 32$ where F is the Fahrenheit temperature (in degrees) and C is the Celsius temperature. What is the Celsius temperature (to the nearest degree) when Fahrenheit temperature is 77° ? 20. _____